

ADDENDUM TO: ON THE RATIONAL APPROXIMATION OF THE SUM OF THE RECIPROCAL OF THE FERMAT NUMBERS

MICHAEL COONS

ABSTRACT. As a corollary of the main result of our recent paper, *On the rational approximation of the sum of the reciprocals of the Fermat numbers* published in this same journal, we prove that for each integer $b \geq 2$ the irrationality exponent of $\sum_{n \geq 0} s_2(n)/b^n$ is equal to 2.

1. INTRODUCTION

Let α be a real number. The *irrationality exponent* $\mu(\alpha)$ is defined as the supremum of the set of real numbers μ such that the inequality

$$\left| \alpha - \frac{p}{q} \right| < \frac{1}{q^\mu}$$

has infinitely many solutions $(p, q) \in \mathbb{Z} \times \mathbb{N}$. For example, Liouville [4] proved that $\mu(\sum_{n \geq 0} 10^{-n!}) = \infty$, and Roth [5] showed that if α is an irrational algebraic number, then $\mu(\alpha) = 2$. Note also that $\mu(\alpha) \geq 2$ for all irrational α .

In this Addendum, we prove the following theorem.

Theorem 1.1. *Let $s_2(n)$ be the sum of the binary digits of n . Then for each integer $b \geq 2$ we have*

$$\mu \left(\sum_{n \geq 0} \frac{s_2(n)}{b^n} \right) = 2.$$

Transcendence of these numbers was proved by Toshimitsu [7] using Mahler's method.

2. PRELIMINARIES

As above, let $s_2(n)$ denote the sum of the binary digits of n , and set $\mathcal{S}(x) := \sum_{n \geq 0} s_2(n)x^n$. Note that for all $n \geq 0$, we have that both $s_2(2n) = s_2(n)$ and $s_2(2n+1) = s_2(n) + 1$. We prove our result by exploiting a connection between the sequences $\{s_2(n)\}_{n \geq 0}$ and $\{f(n)\}_{n \geq 1}$, where we define $f(n)$ by its generating series

$$\mathcal{F}(x) := \sum_{n \geq 1} f(n)x^n = \sum_{n=0}^{\infty} \frac{x^{2^n}}{1+x^{2^n}}.$$

The series $\mathcal{F}(x)$ and its special values have been studied by many authors, including Golomb [3], Duverney [2], and Schwarz [6]. This series is of special interest as $\mathcal{F}(1/2)$ is the sum of the reciprocals of the Fermat numbers. Our interest here is tied to the following result.

Proposition 2.1 (Coons [1]). *Let $b \geq 2$ be a positive integer. Then $\mu(\mathcal{F}(1/b)) = 2$.*

To use Proposition 2.1 to prove Theorem 1.1, we will use the relationship contained in the following lemma.

Lemma 2.2. *For all $n \geq 1$ we have $f(n) = s_2(n) - s_2(n-1)$.*

Date: November 26, 2015.

The research of M. Coons was supported by ARC grant DE140100223.

Proof. Note that the generating function for $\mathcal{F}(x)$ implies that $f(n)$ is multiplicative, and on prime powers given by

$$f(p^k) = \begin{cases} 1 - k & \text{if } p = 2 \\ 1 & \text{if } p \neq 2. \end{cases}$$

Consider the function $v(n) := s_2(n) - s_2(n-1)$ for $n \geq 1$.

To this end, note that if n is odd, say $n = 2k + 1$, then

$$v(2k + 1) = s_2(2k + 1) - s_2(2k) = s_2(k) + 1 - s_2(k) = 1 = f(2k + 1).$$

If n is even, say $n = 2^k(2\ell + 1)$, then

$$\begin{aligned} v(2^k(2\ell + 1)) &= s_2(2^k(2\ell + 1)) - s_2(2^k(2\ell + 1) - 1) \\ &= s_2(2\ell + 1) - s_2(2^{k+1}\ell + 2^k - 1) \\ &= s_2(\ell) + 1 - s_2(2^{k+1}\ell) - s_2(2^k - 1) \\ &= s_2(\ell) + 1 - s_2(\ell) - k \\ &= 1 - k. \end{aligned}$$

From here it is easy to see that $v(n)$ is multiplicative and $f(p^k) = v(p^k)$ for all primes p and integers $k \geq 1$. Thus $v(n) = f(n)$. $\square$

3. PROOF OF THE MAIN RESULT

Proof of Theorem 1.1. Using the above lemma and the fact that $s_2(0) = 0$, we have that

$$\begin{aligned} \mathcal{F}(x) &= \sum_{n \geq 1} f(n)x^n = \sum_{n \geq 1} (s_2(n) - s_2(n-1))x^n = \sum_{n \geq 1} s_2(n)x^n - \sum_{n \geq 1} s_2(n-1)x^n \\ &= \sum_{n \geq 0} s_2(n)x^n - x \sum_{n \geq 0} s_2(n)x^n = (1-x)\mathcal{S}(x), \end{aligned}$$

so that $\mathcal{S}(1/b) = \frac{b}{b-1} \cdot \mathcal{F}(1/b)$. Since $\mathcal{S}(1/b)$ is a (nonzero) rational multiple of $\mathcal{F}(1/b)$, they have the same irrationality exponent. Appealing to Theorem 2.1 proves the theorem. $\square$

Remark. Since $\mathcal{F}(1/b)$ is a (nonzero) rational multiple of $\mathcal{S}(1/b)$, the transcendence of $\mathcal{S}(1/b)$ as proved by Toshimitsu [7] provides an alternative proof (to that of Duverney [2]) of the transcendence of $\mathcal{F}(1/b)$.

REFERENCES

1. Michael Coons, *On the rational approximation of the sum of the reciprocals of the Fermat numbers*, Ramanujan J. **30** (2013), no. 1, 39–65.
2. D. Duverney, *Transcendence of a fast converging series of rational numbers*, Math. Proc. Cambridge Philos. Soc. **130** (2001), no. 2, 193–207.
3. S. W. Golomb, *On the sum of the reciprocals of the Fermat numbers and related irrationalities*, Canad. J. Math. **15** (1963), 475–478.
4. J. Liouville, *Sur des classes très étendues de quantités dont la valeur n'est ni algébrique, ni même réductible à des irrationnelles algébriques*, C.R. Acad. Sci. Paris **18** (1844), 883–885, 910–911.
5. K. F. Roth, *Rational approximations to algebraic numbers*, Mathematika **2** (1955), 1–20; corrigendum, 168.
6. W. Schwarz, *Remarks on the irrationality and transcendence of certain series*, Math. Scand **20** (1967), 269–274.
7. Takeshi Toshimitsu, *Strongly q -additive functions and algebraic independence*, Tokyo J. Math. **21** (1998), no. 1, 107–113.

SCHOOL OF MATHEMATICAL AND PHYSICAL SCIENCES, UNIVERSITY OF NEWCASTLE, AUSTRALIA
E-mail address: Michael.Coons@newcastle.edu.au